


## ATLANTIC Wauseon x Lenape



Suitable for processing – crisps. Oval, yellow tubers with cream flesh. Moderate yields with low outgrade levels, mainly mechanical damage and greens, high dry matter.. Susceptible to splitting, bruising, late blight on foliage and tubers, potato virus Yo, potato cyst nematode *Globodera pallida* Ps2/3,1, gangrene, virus Y and spraing. Resistant to potato cyst nematode *Globodera rostochiensis* Ro1.

Tuber characteristics		Botanical description	
Smoothness of skin	Medium	Maturity	Early Maincrop
Shape of tuber	Short - oval	Height of plants	Medium
Depth of eyes	Shallow - medium	Frequency of berries	Medium
Colour of skin	Cream	Colour of flowers	Red violet
Colour of flesh	Cream	Colour of sprout	Pink

Resistance to Damage, Pests and Diseases	Low	.	.	.	.	.	.	.	High
Late blight on foliage ( <i>Phytophthora infestans</i> )	.	2	3						
Late blight on tubers ( <i>Phytophthora infestans</i> )	.	.	3						
Blackleg ( <i>Pectobacterium atrosepticum</i> )	.	.	.	.	5				
Common scab ( <i>Streptomyces scabiei</i> )	.	.	.	.	.	6			
Potato Cyst Nematode ( <i>Globodera pallida</i> Pa 2/3, 1)	.	2							
Potato Cyst Nematode ( <i>Globodera rostochiensis</i> Ro1)	.	.	.	.	.	.	.	.	9
Potato Leafroll Virus	.	.	.	.	.	6			
Potato Virus Yo	.	.	3						
Bruising	.	.	3						
Splitting	.	.	3						