

SATURNA Maritta x (Record x 1673-1)



| | |
|--|-------------------------------|
| Maturity | = Medium early to medium late |
| Dormancy | = Long |
| Foliage development | = Good to fairly good |
| Colour of skin | = Yellow |
| Colour of flesh | = Pale yellow |
| Shape of tubers | = Round-oval |
| Shallowness of eyes | = Rather deep |
| Size of tubers | = Medium-sized |
| Yield | = Good |
| Dry matter content | = Very high |
| Cooking type | = Fairly firm to floury |
| Suitable for | = Crisps |
| Resistance against leafroll | = Fairly susceptible |
| Resistance against virus A | = Resistant |
| Resistance against virus X | = Fairly good resistance |
| Resistance against virus Yn | = Moderately resistant |
| Resistance against leaf blight | = Moderately resistant |
| Resistance against tuber blight | = Fairly good resistance |
| Wart disease (fysio 1) | = Resistant |
| Resistance to potato cyst nematode Ro1 | = Resistant |
| Common scab | = Fairly susceptible |
| Resistance against internal bruising | = Moderately resistant |



Morphological characteristics

- Plant** tall and intermediate type foliage structure; stems upright to semi-upright and medium to strong anthocyanin colouration; medium, medium to dark green leaves and intermediate to open silhouette; high frequency of flowers, none or very weak anthocyanin colouration inner side flower corolla.
- Tuber** round-oval; yellow, smooth to medium smooth skin; pale yellow flesh; rather deep eyes.
- Sprouts** medium to large, ovoid, strong to very strong intensity of anthocyanin colouration and medium to strong pubescence of base; medium terminal bud with medium anthocyanin colouration of tip; medium root tips.